## SOLAR OBSERVATIONS

## SOLAR AND SKY RADIATION MEASUREMENTS DURING MAY, 1926

By HERBERT H. KIMBALL, Solar Radiation Investigations

For a description of instruments and exposures and an account of the method of obtaining and reducing the measurements, the reader is referred to the Review for January, 1924, 52:42, January, 1925, 53:29, and July, 1925, 53:318.

From Table 1 it is seen that solar radiation intensities averaged slightly above the normal for May at Washington and below the normal at Madison and Lincoln.

At Madison a mid-day intensity of 1.48 gr. cal./min./cm² on the 12th approximates closely to the extreme intensity for May of 1.49 recorded on May 9, 1923.

Table 2 shows an excess in the amount of the radiation received on a horizontal surface from the sun and sky at all three stations, which is most pronounced at Washington.

Skylight polarization measurements made on seven days at Washington give a mean of 60 per cent, with a maximum of 65 per cent on the 27th and 29th. These are higher than the corresponding averages for May at Washington. Measurements made on five days at Madison give a mean of 53 per cent, with a maximum of 63 per cent on the 11th. These are below the corresponding averages for May at Madison.

Table 1.—Solar radiation intensities during May, 1926
[Gram-calories per minute per square centimeter of normal surface]
WASHINGTON, D. C.

	Sun's zenith distance										
	8 a.m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noor
Date	75th mer. time	Air mass									
		A. M.						Р.	м.		solar time
		5.0	4.0	3.0	2.0	*1.0	2.0	3.0	4.0	5.0	e.
Лау 1	mm. 8.48	cal.	cal.	cal.	cal. 0.89	cal.	cal.	cal.	cal.	cal.	mm. 9.47
4 5	3. 99 2. 74 9. 83	0. 64 0. 75	0. 77 0. 84 0. 50		1. 18	1. 40	1.06 1.00			<b>-</b> -	3. 99 2. 89 9 14
8 11 17	7. 57 5. 36 7. 87	0. 49	0.60 0.81 0.74	0.75 0.98		1. 40	1.04	0. 80	0. 67	0. 56	7.0
18 19	10. 97 14. 10	0. 65	0. 74			1. 27	1.00				10. 2 14. 6 7. 8
25 26	6. 27 8. 18 6. 02					1. 27 1. 28	0. 92				4. 7 5. 5
27 29	6. 27 7. 87		0. 60		0. 99	1. 29			(0.45)	(0.54)	6. 2 5. 1
leans Departures			0.72 +0.01	0.86 +0.05	+0.05	1,28 ±0,00		+0.07	(0,67) -0,02		

<sup>•</sup> Extropolated.

TABLE 1.—Solar radiation intensities during May, 1926—Contd.
MADISON. WIS.

	Sun's zenith distance										
	8 a.m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noor
Date	75th mer. time	Air mass									
		A. M.						Ρ.	м.		solar time
		5.0	4.0	3.0	2.0	*1.0	2.0	3.0	4.0	5.0	e.
Ma'y 1	mm. 5, 56	cal.	cal.	cal. 0.69	cal. 0. 91	cal. 1. 29		cal.	cal.	cal.	mm 6.70
5 6 10	6. 02 6. 50				1. 03	1. 33 1. 33 1. 38					9. 8: 6. 0: 5. 7
11 11 12	5. 16			1. 09	1. 24	1. 54					6. 2 5. 7
15 17	7. 29				1. 10	1. 34 1. 28					5. 7 7. 5
20 22	5. 16					1, 22 1, 24					6. 7 5. 3 6. 7
27 28		 			1. 03	1. 39 1. 32					8.1
Means Departures		   <b></b> 		(0, 89) -0, 00		1. 33 0, 02					
			LI	NCO	LN, N	EBR.					
May 3				0. 6		1. 24		0.78	0. 64	0. 5	
5 14	5. 56	0. 7		0.9							10. 2
15 19 24	8. 18	3[	0. 9	1, 0 2 0, 5 0, 9	4 0.77	l	0.9				6. 2 18. 1
Means Departures		(0, 70 -0, 0								(0. 53) -0. 19	

<sup>\*</sup>Extrapolated.

Table 2.—Solar and sky radiation received on a horizontal surface [Gram-calories per square centimeter of horizontal surface]

		Average	A verage daily departure from normal					
Week beginning—	Wash- ington	Madi- son	Lin- coln	Chi- cago	New York	Wash- ington	Madi- son	Lin- coln
1926 Apr. 30	cal. 498	cal. 506	cal. 581	cal. 433	cal. 438	cal. +49	cal. +60	cal. +109
May 7	552	494	318	423	481	+92	+33	-173
21	489 568	465 534	591 630	383 329	350 500	+21 +90	-10 +57	+81 +121
28	532	i 537	564	461	385	+44	+54	+41
Excess since first o	f year on	June 3				+2,765	+2,758	+574

## CORRECTION TO TABLE 2 FOR JANUARY AND FEBRU-ARY, 1926

Owing to a misunderstanding the sensitivity of the recorder in use at Chicago was changed while undergoing repairs in December, 1925. In consequence the daily average for each week for that station published in the Monthly Weather Review for January and February, 1926, should be multiplied by 1.75.